

Unclassified

ENV/JM/MONO(2011)21

Organisation de Coopération et de Développement Économiques
Organisation for Economic Co-operation and Development

27-Jul-2011

English - Or. English

ENV/JM/MONO(2011)21
Unclassified

**GUIDANCE DOCUMENT 117 ON THE CURRENT IMPLEMENTATION OF INTERNAL TRIGGERS
IN TEST GUIDELINE 443 FOR AN EXTENDED ONE GENERATION REPRODUCTIVE TOXICITY
STUDY, IN THE UNITED STATES AND CANADA**

Series on Testing and Assessment

No. 117

JT03305493

Document complet disponible sur OLIS dans son format d'origine
Complete document available on OLIS in its original format

English - Or. English

OECD Environment, Health and Safety Publications

Series on Testing and Assessment

No. 117

**Guidance Document 117 on the Current Implementation of Internal Triggers in Test Guideline 443
for an Extended One Generation Reproductive Toxicity Study, in the United States and Canada**

IOMC

INTER-ORGANIZATION PROGRAMME FOR THE SOUND MANAGEMENT OF CHEMICALS

A cooperative agreement among **FAO, ILO, UNEP, UNIDO, UNITAR, WHO, World Bank and OECD**

Environment Directorate

ORGANISATION FOR ECONOMIC CO-OPERATION AND DEVELOPMENT

Paris 2011

Also published in the Series on Testing and Assessment:

- No. 1, *Guidance Document for the Development of OECD Guidelines for Testing of Chemicals (1993; reformatted 1995, revised 2006)*
- No. 2, *Detailed Review Paper on Biodegradability Testing (1995)*
- No. 3, *Guidance Document for Aquatic Effects Assessment (1995)*
- No. 4, *Report of the OECD Workshop on Environmental Hazard/Risk Assessment (1995)*
- No. 5, *Report of the SETAC/OECD Workshop on Avian Toxicity Testing (1996)*
- No. 6, *Report of the Final Ring-test of the Daphnia magna Reproduction Test (1997)*
- No. 7, *Guidance Document on Direct Phototransformation of Chemicals in Water (1997)*
- No. 8, *Report of the OECD Workshop on Sharing Information about New Industrial Chemicals Assessment (1997)*
- No. 9, *Guidance Document for the Conduct of Studies of Occupational Exposure to Pesticides during Agricultural Application (1997)*
- No. 10, *Report of the OECD Workshop on Statistical Analysis of Aquatic Toxicity Data (1998)*
- No. 11, *Detailed Review Paper on Aquatic Testing Methods for Pesticides and industrial Chemicals (1998)*
- No. 12, *Detailed Review Document on Classification Systems for Germ Cell Mutagenicity in OECD Member Countries (1998)*
- No. 13, *Detailed Review Document on Classification Systems for Sensitising Substances in OECD Member Countries 1998)*
- No. 14, *Detailed Review Document on Classification Systems for Eye Irritation/Corrosion in OECD Member Countries (1998)*
- No. 15, *Detailed Review Document on Classification Systems for Reproductive Toxicity in OECD Member Countries (1998)*
- No. 16, *Detailed Review Document on Classification Systems for Skin Irritation/Corrosion in OECD Member Countries (1998)*

- No. 17, *Environmental Exposure Assessment Strategies for Existing Industrial Chemicals in OECD Member Countries (1999)*
- No. 18, *Report of the OECD Workshop on Improving the Use of Monitoring Data in the Exposure Assessment of Industrial Chemicals (2000)*
- No. 19, *Guidance Document on the Recognition, Assessment and Use of Clinical Signs as Humane Endpoints for Experimental Animals used in Safety Evaluation (1999)*
- No. 20, *Revised Draft Guidance Document for Neurotoxicity Testing (2004)*
- No. 21, *Detailed Review Paper: Appraisal of Test Methods for Sex Hormone Disrupting Chemicals (2000)*
- No. 22, *Guidance Document for the Performance of Out-door Monolith Lysimeter Studies (2000)*
- No. 23, *Guidance Document on Aquatic Toxicity Testing of Difficult Substances and Mixtures (2000)*
- No. 24, *Guidance Document on Acute Oral Toxicity Testing (2001)*
- No. 25, *Detailed Review Document on Hazard Classification Systems for Specifics Target Organ Systemic Toxicity Repeated Exposure in OECD Member Countries (2001)*
- No. 26, *Revised Analysis of Responses Received from Member Countries to the Questionnaire on Regulatory Acute Toxicity Data Needs (2001)*
- No. 27, *Guidance Document on the Use of the Harmonised System for the Classification of Chemicals which are Hazardous for the Aquatic Environment (2001)*
- No. 28, *Guidance Document for the Conduct of Skin Absorption Studies (2004)*
- No. 29, *Guidance Document on Transformation/Dissolution of Metals and Metal Compounds in Aqueous Media (2001)*
- No. 30, *Detailed Review Document on Hazard Classification Systems for Mixtures (2001)*
- No. 31, *Detailed Review Paper on Non-Genotoxic Carcinogens Detection: The Performance of In-Vitro Cell Transformation Assays (2007)*
- No. 32, *Guidance Notes for Analysis and Evaluation of Repeat-Dose Toxicity Studies (2000)*

- No. 33, *Harmonised Integrated Classification System for Human Health and Environmental Hazards of Chemical Substances and Mixtures (2001)*
- No. 34, *Guidance Document on the Development, Validation and Regulatory Acceptance of New and Updated Internationally Acceptable Test Methods in Hazard Assessment (2005)*
- No. 35, *Guidance notes for analysis and evaluation of chronic toxicity and carcinogenicity studies (2002)*
- No. 36, *Report of the OECD/UNEP Workshop on the use of Multimedia Models for estimating overall Environmental Persistence and long range Transport in the context of PBTS/POPS Assessment (2002)*
- No. 37, *Detailed Review Document on Classification Systems for Substances Which Pose an Aspiration Hazard (2002)*
- No. 38, *Detailed Background Review of the Uterotrophic Assay Summary of the Available Literature in Support of the Project of the OECD Task Force on Endocrine Disrupters Testing and Assessment (EDTA) to Standardise and Validate the Uterotrophic Assay (2003)*
- No. 39, *Guidance Document on Acute Inhalation Toxicity Testing (in preparation)*
- No. 40, *Detailed Review Document on Classification in OECD Member Countries of Substances and Mixtures Which Cause Respiratory Tract Irritation and Corrosion (2003)*
- No. 41, *Detailed Review Document on Classification in OECD Member Countries of Substances and Mixtures which in Contact with Water Release Toxic Gases (2003)*
- No. 42, *Guidance Document on Reporting Summary Information on Environmental, Occupational and Consumer Exposure (2003)*
- No. 43, *Guidance Document on Mammalian Reproductive Toxicity Testing and Assessment (2008)*
- No. 44, *Description of Selected Key Generic Terms Used in Chemical Hazard/Risk Assessment (2003)*
- No. 45, *Guidance Document on the Use of Multimedia Models for Estimating Overall Environmental Persistence and Long-range Transport (2004)*
- No. 46, *Detailed Review Paper on Amphibian Metamorphosis Assay for the Detection of Thyroid Active Substances (2004)*

- No. 47, *Detailed Review Paper on Fish Screening Assays for the Detection of Endocrine Active Substances (2004)*
- No. 48, *New Chemical Assessment Comparisons and Implications for Work Sharing (2004)*
- No. 49, *Report from the Expert Group on (Quantitative) Structure-Activity Relationships [(Q)SARs] on the Principles for the Validation of (Q)SARs (2004)*
- No. 50, *Report of the OECD/IPCS Workshop on Toxicogenomics (2005)*
- No. 51, *Approaches to Exposure Assessment in OECD Member Countries: Report from the Policy Dialogue on Exposure Assessment in June 2005 (2006)*
- No. 52, *Comparison of emission estimation methods used in Pollutant Release and Transfer Registers (PRTRs) and Emission Scenario Documents (ESDs): Case study of pulp and paper and textile sectors (2006)*
- No. 53, *Guidance Document on Simulated Freshwater Lentic Field Tests (Outdoor Microcosms and Mesocosms) (2006)*
- No. 54, *Current Approaches in the Statistical Analysis of Ecotoxicity Data: A Guidance to Application (2006)*
- No. 55, *Detailed Review Paper on Aquatic Arthropods in Life Cycle Toxicity Tests with an Emphasis on Developmental, Reproductive and Endocrine Disruptive Effects (2006)*
- No. 56, *Guidance Document on the Breakdown of Organic Matter in Litter Bags (2006)*
- No. 57, *Detailed Review Paper on Thyroid Hormone Disruption Assays (2006)*
- No. 58, *Report on the Regulatory Uses and Applications in OECD Member Countries of (Quantitative) Structure-Activity Relationship [(Q)SAR] Models in the Assessment of New and Existing Chemicals (2006)*
- No. 59, *Report of the Validation of the Updated Test Guideline 407: Repeat Dose 28-Day Oral Toxicity Study in Laboratory Rats (2006)*
- No. 60, *Report of the Initial Work Towards the Validation of the 21-Day Fish Screening Assay for the Detection of Endocrine Active Substances (Phase 1A) (2006)*

- No. 61, *Report of the Validation of the 21-Day Fish Screening Assay for the Detection of Endocrine Active Substances (Phase 1B) (2006)*
- No. 62, *Final OECD Report of the Initial Work Towards the Validation of the Rat Hershberger Assay: Phase-1, Androgenic Response to Testosterone Propionate, and Anti-Androgenic Effects of Flutamide (2006)*
- No. 63, *Guidance Document on the Definition of Residue (2006)*
- No. 64, *Guidance Document on Overview of Residue Chemistry Studies (2006)*
- No. 65, *OECD Report of the Initial Work Towards the Validation of the Rodent Uterotrophic Assay - Phase 1 (2006)*
- No. 66, *OECD Report of the Validation of the Rodent Uterotrophic Bioassay: Phase 2. Testing of Potent and Weak Oestrogen Agonists by Multiple Laboratories (2006)*
- No. 67, *Additional data supporting the Test Guideline on the Uterotrophic Bioassay in rodents (2007)*
- No. 68, *Summary Report of the Uterotrophic Bioassay Peer Review Panel, including Agreement of the Working Group of the National Coordinators of the Test Guidelines Programme on the follow up of this report (2006)*
- No. 69, *Guidance Document on the Validation of (Quantitative) Structure-Activity Relationship [(Q)SAR] Models (2007)*
- No. 70, *Report on the Preparation of GHS Implementation by the OECD Countries (2007)*
- No. 71, *Guidance Document on the Uterotrophic Bioassay - Procedure to Test for Antioestrogenicity (2007)*
- No. 72, *Guidance Document on Pesticide Residue Analytical Methods (2007)*
- No. 73, *Report of the Validation of the Rat Hershberger Assay: Phase 3: Coded Testing of Androgen Agonists, Androgen Antagonists and Negative Reference Chemicals by Multiple Laboratories. Surgical Castrate Model Protocol (2007)*
- No. 74, *Detailed Review Paper for Avian Two-generation Toxicity Testing (2007)*
- No. 75, *Guidance Document on the Honey Bee (Apis Mellifera L.) Brood test Under Semi-field Conditions (2007)*

- No. 76, *Final Report of the Validation of the Amphibian Metamorphosis Assay for the Detection of Thyroid Active Substances: Phase 1 - Optimisation of the Test Protocol (2007)*
- No. 77, *Final Report of the Validation of the Amphibian Metamorphosis Assay: Phase 2 - Multi-chemical Interlaboratory Study (2007)*
- No. 78, *Final Report of the Validation of the 21-day Fish Screening Assay for the Detection of Endocrine Active Substances. Phase 2: Testing Negative Substances (2007)*
- No. 79, *Validation Report of the Full Life-cycle Test with the Harpacticoid Copepods Nitocra Spinipes and Amphiascus Tenuiremis and the Calanoid Copepod Acartia Tonsa - Phase 1 (2007)*
- No. 80, *Guidance on Grouping of Chemicals (2007)*
- No. 81, *Summary Report of the Validation Peer Review for the Updated Test Guideline 407, and Agreement of the Working Group of National Coordinators of the Test Guidelines Programme on the follow-up of this report (2007)*
- No. 82, *Guidance Document on Amphibian Thyroid Histology (2007)*
- No. 83, *Summary Report of the Peer Review Panel on the Stably Transfected Transcriptional Activation Assay for Detecting Estrogenic Activity of Chemicals, and Agreement of the Working Group of the National Coordinators of the Test Guidelines Programme on the Follow-up of this Report (2007)*
- No. 84, *Report on the Workshop on the Application of the GHS Classification Criteria to HPV Chemicals, 5-6 July Bern Switzerland (2007)*
- No. 85, *Report of the Validation Peer Review for the Hershberger Bioassay, and Agreement of the Working Group of the National Coordinators of the Test Guidelines Programme on the Follow-up of this Report (2007)*
- No. 86, *Report of the OECD Validation of the Rodent Hershberger Bioassay: Phase 2: Testing of Androgen Agonists, Androgen Antagonists and a 5 α -Reductase Inhibitor in Dose Response Studies by Multiple Laboratories (2008)*
- No. 87, *Report of the Ring Test and Statistical Analysis of Performance of the Guidance on Transformation/Dissolution of Metals and Metal Compounds in Aqueous Media (Transformation/ Dissolution Protocol) (2008)*

- No.88, *Workshop on Integrated Approaches to Testing and Assessment (2008)*
- No.89, *Retrospective Performance Assessment of the Test Guideline 426 on Developmental Neurotoxicity (2008)*
- No.90, *Background Review Document on the Rodent Hershberger Bioassay (2008)*
- No.91, *Report of the Validation of the Amphibian Metamorphosis Assay (Phase 3) (2008)*
- No.92, *Report of the Validation Peer Review for the Amphibian Metamorphosis Assay and Agreement of the Working Group of the National Coordinators of the Test Guidelines Programme on the Follow-Up of this Report (2008)*
- No.93, *Report of the Validation of an Enhancement of OECD TG 211: Daphnia Magna Reproduction Test (2008)*
- No.94, *Report of the Validation Peer Review for the 21-Day Fish Endocrine Screening Assay and Agreement of the Working Group of the National Coordinators of the Test Guidelines Programme on the Follow-up of this Report (2008)*
- No.95, *Detailed Review Paper on Fish Life-Cycle Tests (2008)*
- No.96, *Guidance Document on Magnitude of Pesticide Residues in Processed Commodities (2008)*
- No.97, *Detailed Review Paper on the use of Metabolising Systems for In Vitro Testing of Endocrine Disruptors (2008)*
- No. 98, *Considerations Regarding Applicability of the Guidance on Transformation/Dissolution of Metals Compounds in Aqueous Media (Transformation/Dissolution Protocol) (2008)*
- No. 99, *Comparison between OECD Test Guidelines and ISO Standards in the Areas of Ecotoxicology and Health Effects (2008)*
- No.100, *Report of the Second Survey on Available Omics Tools (2009)*
- No.101, *Report of the Workshop on Structural Alerts for the OECD (Q)SAR Application Toolbox, 15-16 May 2008, Utrecht, the Netherlands (2009)*
- No. 102, *Guidance Document for using the OECD (Q)SAR Application Toolbox to Develop Chemical Categories According to the OECD Guidance on Grouping of Chemicals (2009)*

- No. 103, *Detailed Review Paper on Transgenic Rodent Mutation Assays (2009)*
- No. 104, *Performance Assessment: Comparison of 403 and CxT Protocols via Simulation and for Selected Real Data Sets (2009)*
- No. 105, *Report on Biostatistical Performance Assessment of the draft TG 436 Acute Toxic Class Testing Method for Acute Inhalation Toxicity (2009)*
- No. 106, *Guidance Document for Histologic Evaluation of Endocrine and Reproductive Test in Rodents (2009)*
- No. 107, *Preservative treated wood to the environment for wood held in storage after treatment and for wooden commodities that are not cover and are not in contact with ground. (2009)*
- No. 108, *Report of the validation of the Hershberger Bioassay (weanling model) (2009)*
- No. 109, *Literature review on the 21-Day Fish Assay and the Fish Short-Term Reproduction Assay (2009)*
- No. 110, *Report of the validation peer review for the weanling Hershberger Bioassay and agreement of the working of national coordinators of the test guidelines programme on the follow-up of this report (2009)*
- No. 111, *Report of the Expert Consultation to Evaluate an Estrogen Receptor Binding Affinity Model for Hazard Identification (2009)*
- No. 112, *The 2007 OECD List of High Production Volume Chemicals (2009)*
- No. 113, *Report of The Focus Session on Current and Forthcoming Approaches for Chemical Safety and Animal Welfare (2010)*
- No. 114, *Performance Assessment of Different Cytotoxic and Cytostatic Measures for the In Vitro Micronucleus Test (MNVT): Summary of results in the collaborative trial (2010)*
- No. 115, *Guidance Document on the Weanling Hershberger Bioassay in Rats: A Short-term Screening Assay for (Anti) Androgenic Properties (2009)*
- No. 116, *Guidance Document on the Design and Conduct of Chronic Toxicity and Carcinogenicity Studies, Supporting TG 451, 452 and 453 (2010)*
- No. 117, *Guidance Document 117 on the Current Implementation of Internal Triggers in Test Guideline 443 for an Extended One*

Generation Reproductive Toxicity Study, in the United States and Canada (2011)

No. 118, *Workshop Report on OECD Countries Activities Regarding Testing, Assessment and Management of Endocrine Disrupters Part I and Part II (2010)*

No. 119, *Classification and Labelling of chemicals according to the UN Globally Harmonized System: Outcome of the Analysis of Classification of Selected Chemicals listed in Annex III of the Rotterdam Convention (2010)*

No. 120, *Part 1: Report of the Expert Consultation on Scientific and Regulatory Evaluation of Organic Chemistry Mechanism-based Structural Alerts for the Identification of DNA Binding Chemicals*

No. 120, *Part 2: Report of the Expert Consultation on Scientific and Regulatory Evaluation of Organic Chemistry Mechanism-based Structural Alerts for the Identification of DNA Binding Chemicals*

No. 121, *Detailed review paper (DRP) on Molluscs life-cycle Toxicity Testing (2010)*

No. 122, *Guidance Document on the determination of the Toxicity of a Test Chemical to the Dung Beetle *Aphodius Constans* (2010)*

No. 123, *Guidance Document on the Diagnosis of Endocrine-related Histopathology in Fish Gonads (2010)*

No. 124, *Guidance for the Derivation of an Acute Reference Dose (2010)*

No. 125, *Guidance Document on Histopathology for Inhalation Toxicity Studies, Supporting TG 412 (Subacute Inhalation Toxicity: 28-Day) and TG 413 (Subchronic Inhalation Toxicity: 90-Day) (2010)*

No. 126, *Short Guidance on the Threshold approach for Acute Fish Toxicity (2010)*

No. 127, *Peer review report of the validation of the 21-day androgenised female stickleback screening assay (2010)*

No. 128, *Validation Report of the 21-day Androgenised Female Stickleback Screening Assay (2010)*

No. 129, *Guidance Document on using Cytotoxicity Tests to Estimate Starting Doses for Acute Oral Systemic Toxicity Tests*

No. 131, *Report of the Test Method Validation of Avian Acute Oral Toxicity Test (OECD test guideline 223) (2010)*

- No. 132, *Report of the Multi-Laboratory Validation of the H295R Steroidogenesis Assay to Identify Modulators (2010)*
- No.133, *Peer Review Report for the H295R Cell-Based Assay for Steroidogenesis (2010)*
- No.134, *Report of the Validation of a Soil Bioaccumulation Test with Terrestrial Oligochaetes by an International ring test (2010)*
- No.135, *Detailed Review Paper on Environmental Endocrine Disruptor Screening: The use of Estrogen and Androgen Receptor Binding and Transactivation Assays in Fish (2010)*
- No.136, *Validation Report of the Chironomid Full Life-Cycle Toxicity Test (2010)*
- No. 137, *Explanatory Background Document to the OECD Test Guideline On In Vitro Skin Irritation Testing (2010)*
- No. 138, *Report of the Workshop on Using Mechanistic Information in Forming Chemical Categories (2011)*
- No. 139, *Report of the Expert Consultation on Scientific and Regulatory Evaluation of Organic Chemistry Mechanism Based Structural Alerts for the Identification of Protein-binding Chemicals (2011)*
- No. 141, *Report of the Phase 1 of the Validation of the Fish Sexual Development Test for the Detection of Endocrine Active Substances (2011)*
- No. 142, *Report of the Phase 2 of the Validation of the Fish Sexual Development Test for the Detection of Endocrine Active Substances (2011)*
- No. 143, *Peer Review Report for the Validation of the Fish Sexual Development Test and Agreement of the Working Group of National Co-ordinators of the Test Guideline Programme on the Follow-up of the Peer Review (2011)*
- No. 144, *Validation report supporting the Test Guideline for the acute chironomid assay (2011)*

No. 145, *Retrospective Performance Assessment supporting the Development of the Test Guideline on Transgenic Rodent Somatic and Germ Cells Gene Mutation Assays (2011)*

© **OECD 2011**

Applications for permission to reproduce or translate all or part of this material should be made to: Head of Publications Service,
RIGHTS@oecd.org. OECD, 2 rue André-Pascal, 75775 Paris Cedex
16, France

ABOUT THE OECD

The Organisation for Economic Co-operation and Development (OECD) is an intergovernmental organisation in which representatives of 34 industrialised countries in North and South America, Europe and the Asia and Pacific region, as well as the European Commission, meet to co-ordinate and harmonise policies, discuss issues of mutual concern, and work together to respond to international problems. Most of the OECD's work is carried out by more than 200 specialised committees and working groups composed of member country delegates. Observers from several countries with special status at the OECD, and from interested international organisations, attend many of the OECD's workshops and other meetings. Committees and working groups are served by the OECD Secretariat, located in Paris, France, which is organised into directorates and divisions.

The Environment, Health and Safety Division publishes free-of-charge documents in ten different series: Testing and Assessment; Good Laboratory Practice and Compliance Monitoring; Pesticides and Biocides; Risk Management; Harmonisation of Regulatory Oversight in Biotechnology; Safety of Novel Foods and Feeds; Chemical Accidents; Pollutant Release and Transfer Registers; Emission Scenario Documents; and Safety of Manufactured Nanomaterials. More information about the Environment, Health and Safety Programme and EHS publications is available on the OECD's World Wide Web site (www.oecd.org/ehs/).

This publication was developed in the IOMC context. The contents do not necessarily reflect the views or stated policies of individual IOMC Participating Organisations.

The Inter-Organisation Programme for the Sound Management of Chemicals (IOMC) was established in 1995 following recommendations made by the 1992 UN Conference on Environment and Development to strengthen co-operation and increase international co-ordination in the field of chemical safety. The Participating Organisations are FAO, ILO, UNEP, UNIDO, UNITAR, WHO, World Bank and OECD. UNDP is an observer. The purpose of the IOMC is to promote co-ordination of the policies and activities pursued by the Participating Organisations, jointly or separately, to achieve the sound management of chemicals in relation to human health and the environment.

This publication is available electronically, at no charge.

**For this and many other Environment,
Health and Safety publications, consult the OECD's
World Wide Web site (www.oecd.org/ehs/)**

or contact:

**OECD Environment Directorate,
Environment, Health and Safety Division
2 rue André-Pascal
75775 Paris Cedex 16
France**

Fax: (33-1) 44 30 61 80

E-mail: ehscont@oecd.org

FOREWORD

This Guidance Document provides guidance on the current implementation of internal triggers in TG 443: Extended One Generation Reproductive Toxicity Study, in the United States and Canada.

A combined meeting of the National Coordinators of the Test Guidelines Programme and of an expert group on reproductive toxicity studies, held in Arlington on 19-21 October 2010, agreed to remove the text concerning the procedure for the decision on the internal triggers from TG 443. The Arlington meeting and the Joint Meeting of the Chemicals Committee and the working Party on Chemicals, Pesticides and Biotechnology (Joint Meeting), however, agreed to include internal triggers in a Guidance Document that would be published at the same time as TG 443.

The Guidance Document was prepared by the United States and Canada. It was revised and approved by the WNT at its meeting in April 2011. The Joint Meeting agreed to the declassification of this document on 12 July 2011.

This document is published under the responsibility of the Joint Meeting.

Guidance Document 117 on the current implementation of internal triggers in Test Guideline 443 for an Extended One Generation Reproductive toxicity Study, in the Unites States and Canada

1. This Guidance Document has been developed to support the use of Test Guideline 443 for an Extended One-Generation Reproduction Toxicity Study (EOGRTS) under Canadian (Pest Management Regulatory Agency) and US regulatory authorities (United States Environmental Protection Agency, Office of Pesticide Programs). Internal triggers for the assessment of the second generation required by these authorities are discussed in Annex 1. This document is intended to provide guidance to users of TG 443 under these authorities. Other regions or member states can have different requirements or approaches for implementing TG 443. A more general OECD Guidance Document on the EOGRTS is in preparation.
2. Under the Canadian Pest Management Regulatory Agency and the United States Environmental Protection Agency, Office of Pesticide Programs, the decision to assess a second generation will take into consideration all available information including data gathered during the conduct of the EOGRTS. In making this determination, consideration should be given to issues such as the sensitivity of the young noted in the database and seriousness of pre- or post-natal endpoints, although other factors can influence the assessment (i.e., dose-response, toxicokinetics and/or metabolism, mode of action, confidence in study and/or endpoints, and human exposure information). It is expected that, with the additional parameters evaluated in the F₁ generation in the EOGRTS, the F₂ with their limited parameter assessments would seldom affect the hazard characterization for risk assessment (Piersma et al., 2010). However, until sufficient empirical data can be provided to support this hypothesis, the decision by the Canadian Pest Management Regulatory Agency and the United States Environmental Protection Agency, Office of Pesticide Programs to assess the second generation will be determined through the implementation of triggers, as defined in this Guidance Document.
3. As noted above, these triggers are intended to be used for the hazard characterization in risk assessments rather than for Classification and Labelling within the mentioned jurisdictions. Companies wishing to register chemicals in other regions are expected to consult regulatory authorities to determine the appropriate testing protocol.
4. When a maximum of 20 EOGRTS have been conducted with an assessment of the second generation, the available results and appropriate scientific evidence will be analysed. Subsequently, this Guidance Document will be reviewed and, if necessary, revised in light of experience gained. In this regard, OECD member countries and regions are encouraged to share available results.

Annex

Current implementation of TG 443 in the US and Canadian regulatory frameworks for pesticides

Text provided by Canada and USA¹:

Introduction

5. The OECD expert group for the EOGRTS held several meetings to discuss the interpretation of the data derived from Test Guideline 443. The meetings recognized the need for a Guidance Document on triggers for assessing the second generation and on methodological aspects and interpretation of data for all three cohorts within the Test Guideline (reproductive, developmental immunotoxicity, and developmental neurotoxicity).
6. This document constitutes an essential supplement to OECD Test Guideline 443 (EOGRTS) to describe the use of data gathered within the study to determine if a second generation should be produced. However, as a starting point, available mode of action information and data from other toxicity studies (e.g., repeat dose toxicity studies for systemic toxicity, and ADME) should be considered in the assessment as they may already indicate the potential for reproductive toxicity.
7. The EOGRTS design includes evaluations of numerous sensitive structural, functional, and endocrine-mediated components. Thus, it is unlikely that a critical effect on development and reproduction would be missed. Using a science and risk based approach (as described in Cooper et al., 2006) to determine the need for an F₂ evaluation allows for a tailored approach to testing, reduces the numbers of animals used (1200 animals are used to generate an F₂), and the resources needed to manage, review, and document the study. If required, the production of an F₂ (i.e., breeding of the Cohort 1 F_{1b} animals) does represent a critical decision point integral to the study design. This decision will need to be made rapidly with a clear understanding of supporting data. When determining whether production of an F₂ is needed, one should consider how the additional information gained by breeding a second generation will be used in the hazard evaluation or overall risk assessment.
8. Typically, the F₂ generation allows for a replicate assessment of reproductive performance, litter size, offspring survival and development (including anogenital distance and nipple retention), and weanling necropsy endpoints (organ weights and histopathology) after a lifetime of exposure. A weight of evidence will be essential to determine if the relative sensitivity of the young and seriousness of pre- and post-natal endpoints have been sufficiently evaluated for the hazard characterization for risk assessment.
9. Lastly, when predicted human exposures are considered adequately characterized, margin of exposure (MOE) considerations may be factored into the decision to require the assessment of a second generation. For example, if toxicity triggers are limited to the high dose level alone (with no apparent dose-related trend), MOE of this dose relative to either estimated human exposures or those directly measured through human biomonitoring studies could help guide the triggering decision.

¹ This annex is provided for transparency reasons. It represents the views from the US and Canada. It has not been approved by other OECD Member States.

Determining the level of concern

10. Hazard identification involves understanding the inherent properties of a substance that may lead to adverse responses. It involves defining the relationship between the dose of a substance administered to or received by the test species and the qualitative and quantitative response to the substance.
11. When results of animal testing reveal unique effects in the young (e.g. a different pattern of effects of concern) relative to adults, this is referred to as susceptibility or qualitative sensitivity. Evidence of quantitative sensitivity arises when the effects in the young occur at doses lower than those causing effects in adults, occur more quickly, or occur with greater severity or duration than those observed in adults.
12. If the critical endpoint is based on a serious toxicological effect, a high degree of concern would be identified. The temporal nature of the effect (e.g. time of onset, persistence, recovery, etc.) will influence the determination of the degree of concern, with irreversible findings eliciting greater concern. Examples of serious endpoints of concern include, but are not limited to, reduced viability of offspring, and the occurrence of malformations. Less serious endpoints of concern could include transient effects such as organ weight changes with no or minimal associated histopathological changes.
13. As described above, an evaluation of pre- and post-natal toxicity, primarily determined by an analysis of the sensitivity of the young and seriousness of pre- or post-natal endpoints, is critical to the hazard characterization for risk assessment under certain regulatory authorities. A recent retrospective analysis of 498 multi-generation reproductive toxicity studies (OECD TG 416 and USEPA 870.3800), representing 438 different tested substances, determined that the second generation mating and offspring will rarely provide critical information for hazard identification (Piersma et al., 2010). However, until sufficient empirical data can be provided to support this hypothesis in the context of the EOGRTS, under certain regulatory authorities, the assessment of the second generation will be determined through the implementation of triggers, as defined below.

Determination of Triggers for the Assessment of the Second Generation in the EOGRTS

14. Table 1 below provides a list of endpoints (or “triggers”) which can be used to determine whether or not an assessment of the second generation is required. Criteria for establishing triggers were based on the original proposal by Cooper et al. (2006) as well as discussions within the expert group for the EOGRTS.
15. Several general categories of triggers for the production of a second-generation are proposed in order to better characterize equivocal results and/or the relative sensitivity of the young and/or seriousness of pre- and post-natal endpoints:
 - an adverse effect on fertility or fecundity of the parental generation,
 - indications of abnormal sexual development of the F₁ pups,
 - adverse effects on F₁ litter parameters and developmental landmarks,
 - death or evidence of toxicity to the F₁ pups pre-weaning, and
 - equivocal effects on F₁ parameters or unusual control data compared to historical background may also trigger a second generation

Adverse effects on fertility or fecundity of the parental generation

16. There are some endpoints that are only examined in the P animals and/or F₁ offspring (Table 1). These include reproductive performance (number of implantations, pregnancy rate, and gestation interval) and estrous cyclicity. Dose-dependant and biologically relevant alterations in these endpoints justify the production of a second generation, particularly if these findings occur in the absence of severe maternal toxicity. In this situation, evaluating a second generation would provide an assessment of the consequences of these types of effects, after a lifetime of exposure.
17. It is generally accepted that male reproductive organ histopathology is the most sensitive endpoint for detecting minor changes in spermatogenesis in rats (Mangelsdorf *et al.*, 2003; Ulbrich and Palmer, 1995). Functional evaluations of fertility are less sensitive due to the excess sperm reserve in rodents. Mangelsdorf *et al.* (2003), in an assessment for the German Federal Institute on Occupational Safety and Health, reported that reproductive/accessory sex gland organ weights and sperm parameters (motility and counts) were more sensitive endpoints for detecting toxicant effects on reproduction than fertility parameters (number of implantations and pregnancies). These results are consistent with a limited analysis by Gray *et al.* (1989) who reported effects on sperm and gonadal toxicity occurred at lower doses than effects on fertility. These data consistently support the premise that alterations in sperm parameters will be more sensitive at detecting potential adverse effects than the assessment of a second generation.
18. Given the greater sensitivity of histopathology and sperm evaluations to detect changes in male reproductive toxicity compared to the functional observation by the assessment of a second generation, neither effects on reproductive histopathology nor effects on sperm parameters warrant breeding a second generation. These endpoints are assessed twice in the EOGRTS, including an assessment of F₁ offspring that have been exposed *in utero*, during lactation and maturation. F₁ offspring data on reproductive organ weights, histopathology and andrology will not be available when a decision to mate the F₁ adults is needed (Table 1); however, the greater sensitivity of these endpoints fulfills risk assessment needs better than the assessment of a second generation. Furthermore, neither mature testicular histopathology nor sperm assessment is assessed in F₂ animals, which are euthanized at weaning.
19. With respect to the female reproductive endpoints (estrous cycle evaluation, reproductive organ weights and histopathology, and ovarian follicle counts), there are fewer data available for the comparison of endpoint sensitivity. Generally, it is recognized that alterations in ovarian follicle development in female rats may not affect fertility (Hirshfield, 1987). Female rats have robust reproductive performance and hormone production even in the presence of reproductive system alterations (*e.g.*, within 24 hours of removing an ovary, the remaining ovary can ovulate a full complement of 10-12 follicles; only 2-3 corpora lutea are needed to maintain pregnancy – Hirshfield, 1987). Thus, ovarian histopathology is believed to be a sensitive indicator of female reproductive toxicity (Regan *et al.*, 2005). Examination of adult ovarian histopathology is favoured as decreases in primordial follicle numbers will be exacerbated in adults due to continuous recruitment of the remaining follicles (Regan *et al.*, 2005). Therefore, examination of adult P and F₁ offspring is preferred over an examination of prepubescent F₂ weanlings. While the Society of Toxicologic Pathology favours qualitative ovarian histopathology (conducted in the Ps) as an initial assessment of ovarian effects (Regan *et al.*, 2005), Bolon *et al.* (1997) has suggested that ovarian follicle counts (conducted in the F₁) also provide a more sensitive indicator of female reproductive toxicity than fertility, again favouring a histopathology endpoint over fertility.

20. With respect to reproductive organ weights, uterine weight is highly variable, depending on the stage of the estrous cycle at necropsy. The stage of the estrous cycle at the time of necropsy is also not a predictive endpoint, being limited to a single time-point. Effects on reproductive organ weights and histopathology do not warrant the assessment of a second generation as these endpoints are either more, or as, sensitive to toxicant alterations than fertility and because additional information on these endpoints would not be obtained in F₂ pups. A second assessment of reproductive organ weights and histopathology will be available from the F₁ adults, including ovarian follicle counts. These data should fulfil risk assessment needs.

Abnormal sexual development of the F₁ pups, adverse effects on F₁ litter parameters and developmental landmarks, and death or evidence of toxicity to the F₁ pups pre-weaning:

21. Effects on 1) F₁ litter size in the absence of P reproductive organ histopathology changes or 2) pup survival in the absence of severe maternal toxicity or 3) pup developmental landmarks (discussed below) would require the assessment of a second generation in order to determine the relative sensitivity of the young and seriousness of pre- and post-natal endpoints.
22. Puberty is only examined in the F₁ offspring (3/sex/litter), but effects seen in the absence of body weight effects may indicate that the underlying development of the reproductive system has been affected. Some regulatory authorities may therefore require an assessment of the second generation.
23. There are some endpoints that, without producing a second generation, are only examined in the F₁ offspring (Table 1). These include litter size and weight, offspring survival, offspring development (including anogenital distance and nipple retention) and endpoints assessed at weaning necropsy. Treatment- and dose-related alterations in these endpoints in the absence of severe maternal toxicity, justifies the production of a second generation. As with other toxicological studies, weight of evidence will be applied when interpreting the results of the EOGRTS. The weight of evidence concept becomes especially important given the number of endpoints examined and the opportunity for Type I error. The laboratory's historical control data (HCD) can assist in the interpretation of data on reproductive toxicity endpoints.

Equivocal effects on F₁ parameters or unusual control data compared to historical background may also trigger a second generation:

24. As discussed in the Test Guideline 443, the assessment of a second generation may help clarify equivocal findings or provide further characterization of effects on fertility observed in the first generation mating.
25. Statistical significance does not need to be present to validate the biological significance of treatment-related effects. This is particularly true of findings with low incidence (i.e., rare malformations) or high variability, or in situations where the concurrent control data have an unusual incidence profile. In the same way, statistical significance does not necessarily signify biological significance, and scientific judgement and relevant historical control data should be used to distinguish between spurious and real findings (OECD GD 43).

26. Comparison of concurrent study control data with the data from treated animals should always take precedence over comparison with historical control data. The primary goal of evaluating the historical control data is to determine if concurrent control animals are behaving within the margins of normal variability for the species and strain. If historical control data are demonstrably different from concurrent control data, it may be an indication that the study contains some flaw. In the most egregious case, it may not be appropriate to utilise the historical control data in the interpretation of data from treated groups (OECD GD43).

27. If historical control data are used, the most appropriate of these are from studies conducted in the same laboratory, within a reasonable amount of time prior to the study being interpreted (*e.g.*, ± 2 years) in order to avoid genetic drift in the laboratory animal population, and under the same study conditions (*e.g.*, identical species, strain, source, age, vehicle, route and duration of administration, technical personnel, etc.). It is important that the data include sufficient information to render it meaningful in the context of the concurrent study. For example, definitions of terminology should be provided; incidental and continuous data should be fully characterized and summarised with appropriate data ranges, maximum, minimum, median, and mean values; data variance should be addressed. Historical control information that is compiled by animal suppliers or through surveys (or including genetic drift) across multiple laboratories (Clemens *et al.*, 1994, MARTA and MTA, 1995, 1996) can also be useful in some situations and under the appropriate caveats. Overall, the interpretation and use of historical control data requires careful consideration, and the application of scientific judgement and expertise (OECD GD43).

Table 1. Criteria for assessing the second generation in the EOGRTS

Trigger Endpoints ^a	Recommendations
Adult Endpoints	
P Fertility (# implantations, pregnancy rate, gestational interval)	Mate F ₁ in the absence of corresponding biologically relevant and dose-related changes in reproductive histopathology
F ₁ Estrous Cycle Evaluation	Mate F ₁ if biologically relevant and dose-related changes in estrous cycle length without severe toxicity in the dams ^b
Offspring Endpoints	
F ₁ Litter parameters (litter size)	Mate F ₁ if biologically relevant and dose-related decreases in litter size are seen in the absence of severe maternal toxicity or lethality ^b
F ₁ Developmental landmarks (AGD, nipple retention, puberty onset, PPS, VO)	Mate F ₁ if biologically relevant and dose-related effects in the absence of body weight-mediated changes in these parameters
↓ F ₁ pup survival post-natally	Mate F ₁ in the absence of severe maternal toxicity ^b
F ₁ pup malformations	Mate F ₁ in the absence of severe maternal toxicity ^b
↓ F ₁ live birth index	Mate F ₁ in the absence of severe maternal toxicity ^b
↓ F ₁ pup body weight	Mate F ₁ , if pup body weight decrease is biologically relevant and in the absence of maternal body weight decrements

^a Each endpoint will be available in sufficient time to determine whether or not the F₁ should be mated.

^b Type, incidence, magnitude and severity of effect(s) should be considered in relation to maternal toxicity.

REFERENCES

- Bolon, B., Bucci, T. J., Wargritton, A. R., Chen, J. J., Mattison, D. R., and Heindel, J. J. (1997). Differential follicle counts as a screen for chemically induced ovarian toxicity in mice: results from continuous breeding bioassays. *Fundam. Appl. Toxicol.* 39, 1-10.
- Cooper, R. L., Lamb, J. C., Barlow, S. M., Bentley, K., Brady, A. M., Doerr, N., Eisenbrandt, D. L., Fenner-Crisp, P. A., Hines, R. N., Irvine, L. F. H., Kimmel, C. A., Koeter, H., Li, A. A., Makris, S. L., Sheets, L. P., Speijers, G. J. A., and Whitby, K. E. (2006). A tiered approach to life stages testing for agricultural chemical safety assessment. *Crit. Rev. Toxicol.* 36, 69-98.
- Food and Drug Regulations, Section C.01.001., Part C Drugs, Division1, 4 December 1997.
- Gray, L. E., Jr., Ostby, J., Sigmon, R., Ferrell, J., Rehnberg, G., Linder, R., Cooper, R., Goldman, J., and Laskey, J. (1988). The development of a protocol to assess reproductive effects of toxicants in the rat. *Reprod. Toxicol.* 2, 281-287.
- Gray, L., Ostby, J., Ferrell, J., Sigmon, R., Cooper, R., Linder, R., Rehnberg, G., Goldman, J., and Laskey, J. (1989). Correlation of sperm and endocrine measures with reproductive success in rodents. In: *Progress in Clinical and Biological Research*, vol. 302 (E.J. Burger, ed.) Liss, New York, pp. 193-209.
- Hirshfield, A. N. (1987). Histological assessment of follicular development and its applicability to risk assessment. *Reprod. Toxicol.* 1, 71-79.
- Mangelsdorf, I., Buschmann, J., and Orthen, B. (2003). Some aspects relating to the evaluation of the effects of chemicals on male fertility. *Regul. Toxicol. Pharmacol.* 37, 356-369.
- OECD Guidance Document on Mammalian Reproductive Toxicity Testing and Assessment No. 43 (GD 43). ENV/JM/MONO(2008)16.
- Piersma AH, Rorije E, Beekhuijzen ME, Cooper R, Dix DJ, Heinrich-Hirsch B, Martin MT, Mendez E, Muller A, Paparella M, Ramsingh D, Reaves E, Ridgway P, Schenk E, Stachiw L, Ulbrich B, Hakkert BC. (2010). Combined retrospective analysis of 498 rat multi-generation reproductive toxicity studies: On the impact of parameters related to F1 mating and F2 offspring. *Reprod Toxicol* Dec 3.
- Regan, K. S., Cline, J. M., Creasy, D., Davis, B., Foley, G. L., Lanning, L., Latendresse, J. R., Makris, S., Morton, D., Rehm, S., and Stebbins, K. (2005). STP position paper: Ovarian follicle counting in the assessment of rodent reproductive toxicity. *Toxicol. Pathol.* 33, 409-412.
- Ulbrich, B., and Palmer, A. K. (1995). Detection of effects on male reproduction – a literature survey. *J. Am. Coll. Toxicol.* 14 (4), 293-327.